

A REDUCTION MAP FOR NEF LINE BUNDLES

TH. BAUER, F. CAMPANA, TH. ECKL, S. KEBEKUS, TH. PETERNELL,
S. RAMS, T. SZEMBERG, AND L. WOTZLAW

CONTENTS

1. Introduction	1
2. A reduction map for nef line bundles	2
2.1. Construction of the reduction map	2
2.2. Nef cohomology classes	5
2.3. The nef dimension	5
2.4. The structure of the reduction map	6
3. A Counterexample	7
References	9

1. INTRODUCTION

In [Ts00], H. Tsuji stated several very interesting assertions on the structure of pseudo-effective line bundles L on a projective manifold X . In particular he postulated the existence of a meromorphic “reduction map”, which essentially says that through the general point of X there is a maximal irreducible L -flat subvariety. Moreover the reduction map should be almost holomorphic, i.e. has compact fibers which do not meet the indeterminacy locus of the reduction map. The proofs of [Ts00] however are extremely difficult to follow.

The purpose of this note is to establish the existence of a reduction map in the case where L is nef and to prove that it is almost holomorphic —this was also stated explicitly in [Ts00]. Our proof is completely algebraic while [Ts00] works with deep analytic methods. Finally, we show by a basic example that in the case where L is only pseudo-effective, the postulated reduction map cannot be almost holomorphic —in contrast to a claim in [Ts00].

These notes grew out from a small workshop held in Bayreuth in January 2001, supported by the Schwerpunktprogramm “Global methods in complex geometry” of the Deutsche Forschungsgemeinschaft.

2. A REDUCTION MAP FOR NEF LINE BUNDLES

In this section we want to prove the following structure theorem for nef line bundles on a projective variety.

Theorem 2.1. *Let L be a nef line bundle on a normal projective variety X . Then there exists an almost holomorphic, dominant rational map $f : X \dashrightarrow Y$ with connected fibers, called a “reduction map” such that*

1. *L is numerically trivial on all compact fibers F of f with $\dim F = \dim X - \dim Y$*
2. *for every general point $x \in X$ and every irreducible curve C passing through x with $\dim f(C) > 0$, we have $L \cdot C > 0$.*

The map f is unique up to birational equivalence of Y .

This theorem was stated without complete proof in Tsuji’s paper [Ts00]. Relevant definitions are given now.

Definition 2.2. *Let X be an irreducible reduced projective complex space (projective variety, for short). A line bundle L on X is numerically trivial, if $L \cdot C = 0$ for all irreducible curves $C \subset X$. The line bundle L is nef if $L \cdot C \geq 0$ for all curves C .*

Let $f : Y \rightarrow X$ be a surjective map from a projective variety Y . Then clearly L is numerically trivial (nef) if and only if $f^*(L)$ is.

Definition 2.3. *Let X and Y be normal projective varieties and $f : X \dashrightarrow Y$ a rational map and let $X^0 \subset X$ be the maximal open subset where f is holomorphic. The map f is said to be almost holomorphic if some fibers of the restriction $f|_{X^0}$ are compact.*

2.1. Construction of the reduction map.

2.1.1. *A criterion for numerical triviality.* In order to prove theorem 2.1 and construct the reduction map, we will employ the following criterion for a line bundle to be numerically trivial.

Theorem 2.4. *Let X be an irreducible projective variety which is not necessarily normal. Let L be a nef line bundle on X . Then L is numerically trivial if and only if any two points in X can be joined by a (connected) chain C of curves such that $L \cdot C = 0$.*

In the remaining part of the present section 2.1.1 we will prove theorem 2.4. The proof will be performed by a reduction to the surface case. The argumentation is then based on the following statement, which, in the smooth case, is a simple corollary to the Hodge Index Theorem.

Proposition 2.5. *Let S be an irreducible, projective surface which is not necessarily normal, and let $q : S \rightarrow T$ be a morphism with connected fibers onto a curve. Assume that there exists a nef line bundle $L \in \text{Pic}(S)$ and a curve $C \subset S$ such that $q(C) = T$ and such that*

$$L \cdot F = L \cdot C = 0$$

holds, where F is a general q -fiber. Then L is numerically trivial.

Proof. If S is smooth, set $D = C + nF$, where n is a large positive integer. Then we have $D^2 > 0$. By the Hodge Index Theorem it follows that

$$(L \cdot D)^2 \geq L^2 \cdot D^2,$$

hence $L^2 = 0$, since by our assumptions $L \cdot D = 0$. So equality holds in the Index Theorem and therefore L and D are proportional: $L \equiv kD$ for some rational number k . Since $0 = L^2 = k^2 D^2$ and $D^2 > 0$, we conclude that $k = 0$. That ends the proof in the smooth case.

If S is singular, let $\delta : \tilde{S} \rightarrow S$ be a desingularisation of S and let $\tilde{C} \subset \tilde{S}$ be a component of $\delta^{-1}(C)$ which maps surjectively onto C . Note that the fiber of $q \circ \delta$ need no longer be connected and consider the Stein factorisation

$$\begin{array}{ccc} \tilde{S} & \xrightarrow[\text{desing.}]{\delta} & S \\ \tilde{q} \downarrow & & \downarrow q \\ \tilde{T} & \xrightarrow[\text{finite}]{} & T \end{array}$$

It follows immediately from the construction that $\tilde{q}(\tilde{C}) = \tilde{T}$, that $\delta^*(L)$ has degree 0 on $\tilde{C}$ and on the general fiber of $\tilde{q}$. The argumentation above therefore yields that $\delta^*(L)$ is trivial on $\tilde{S}$. The claim follows. $\square$

2.1.2. Proof of Theorem 2.4. If L is numerically trivial, the assertion of theorem 2.4 is clear. We will therefore assume that any two points can be connected by a curve which intersects L with multiplicity 0, and we will show that L is numerically trivial. To this end, choose an arbitrary, irreducible curve $B \subset X$. We are finished if we show that $L \cdot B = 0$.

Let $a \in X$ be an arbitrary point which is not contained in B . For any $b \in B$ we can find by assumption a connected, not necessarily irreducible, curve Z_b containing a and b such that $L \cdot Z_b = 0$. Since the Chow variety has compact components and only a countable number of components, we find a family $(Z_t)_{t \in T}$ of curves, parametrised by a compact irreducible curve $T \subset \text{Chow}(X)$ such that for every point $b \in B$, there exists a point $t \in T$ such that the curve Z_t contains both a and b . We consider the universal family $S \subset X \times T$ over T together with the projection morphisms

$$\begin{array}{ccc} S & \xrightarrow{p} & X \\ q \downarrow & & \\ T & & \end{array}$$

Claim 1. There exists an irreducible component $S_0 \subset S$ such that $p^*(L)$ is numerically trivial on S_0 .

Proof of claim 1. As all curves Z_t contain the point a , the surface S contains the curve $\{a\} \times T$. Let $S_0 \subset S$ be a component which contains $\{a\} \times T$. Since $\{a\} \times T$ intersects all fibers of the natural projection morphism q , and since $p^*(L)$ is trivial on $\{a\} \times T$, an application of proposition 2.5 yields the claim. $\square$

Claim 2. The bundle $p^*(L)$ is numerically trivial on S .

Proof of claim 2. We argue by contradiction and assume that there are components $S_j \subset S$ where $p^*(L)$ is not numerically trivial. We can therefore subdivide S into two subvarieties as follows:

$$\begin{aligned} S_{triv} &:= \{\text{union of the irreducible components } S_i \subset S \\ &\quad \text{where } p^*(L)|_{S_i} \text{ is numerically trivial}\} \\ S_{ntriv} &:= \{\text{union of the irreducible components } S_i \subset S \\ &\quad \text{where } p^*(L)|_{S_i} \text{ is not numerically trivial}\} \end{aligned}$$

By assumption, and by claim 1, both varieties are not empty. Since S is the universal family over a curve in $\text{Chow}(X)$, the morphism q is equidimensional. In particular, since all components $S_i \subset S$ are two-dimensional, every irreducible component S_i maps surjectively onto T . Thus, if $t \in T$ is a general point, the connected fiber $q^{-1}(t)$ intersects both S_{triv} and S_{ntriv} . Thus, over a general point $t \in T$, there exists a point in $S_{triv} \cap S_{ntriv}$. It is therefore possible to find a curve $D \subset S_{triv} \cap S_{ntriv}$ which dominates T .

That, however, contradicts proposition 2.5: on one hand, since $D \subset S_{triv}$, the degree of $p^*(L)|_D$ is 0. On the other hand, we can find an irreducible component $S_j \subset S_{ntriv} \subset S$ which contains D . But because $p^*(L)$ has degree 0 on the fibers of $p^*(L)|_{S_j}$, proposition 2.5 asserts that $p^*(L)$ is numerically trivial on S_j , contrary to our assumption. This ends the proof of Claim 2. $\square$

We apply Claim 2 as follows: if $B' \subset S$ is any component of the preimage $p^{-1}(B)$, then $p^*(L).B' = 0$. That shows that $L.B = 0$, and the proof of theorem 2.4 is done. $\square$

2.1.3. Proof of Theorem 2.1. In order to derive Theorem 2.1 from Theorem 2.4, we introduce an equivalence relation on X with setting $x \sim y$ if x and y can be joined by a connected curve C such that $L \cdot C = 0$. Then by [Ca81] or [Ca94, appendix], there exists an almost holomorphic map $f : X \dashrightarrow Y$ with connected fibers to a normal projective variety Y such that, two general points x and y satisfy $x \sim y$ if and only if $f(x) = f(y)$. This map f gives the fibration we are looking for.

If F is a general fiber, then $L|_F \equiv 0$ by Theorem 2.4.

We still need to verify that $L \cdot C = 0$ for all curves C contained in an arbitrary compact fiber F_0 of dimension $\dim F_0 = \dim X - \dim Y$. To do that, let H be an ample line bundle on X and pick

$$D_1, \dots, D_k \in |mH|$$

for m large such that

$$D_1 \cdot \dots \cdot D_k \cdot F_0 = C + C'$$

with an effective curve C' . Then

$$L \cdot (C + C') = L \cdot D_1 \cdot \dots \cdot D_k \cdot F$$

with a general fiber F of f , hence $L \cdot (C + C') = 0$. Since L is nef, we conclude $L \cdot C = 0$. $\square$

2.2. Nef cohomology classes. In Theorems 2.1 and 2.4 we never really used the fact that L is a line bundle; only the property that $c_1(L)$ is a nef class is important and even rationality of the class does not play any role. Hence our results directly generalize to nef cohomology classes of type $(1,1)$. To be precise, we fix a projective manifold (we stick to the smooth case for sakes of simplicity) and we say that a class $\alpha \in H^{1,1}(X, \mathbb{R})$ is nef, if it is in the closure of the cone generated by the Kähler classes. Moreover α is numerically trivial, if $\alpha \cdot C = 0$ for all curves $C \subset X$.

If $Z \subset X$ is a possibly singular subspace, then we say that α is numerically trivial on Z , if for some (and hence for all, see [Pa98]) desingularisation $\hat{Z} \rightarrow Z$, the induced form $f^*(\alpha)$ is numerically trivial on $\hat{Z}$, i.e. $f^*(\alpha) \cdot C = 0$ for all curves $C \subset \hat{Z}$. Here $f : \hat{Z} \rightarrow X$ denotes the canonical map. Similarly we define α to be nef on Z . If Z is smooth, this is the same as to say that $\alpha|_Z$ is a nef cohomology class in the sense that $\alpha|_Z$ is in the closure of the Kähler cone of Z .

Theorem 2.6. *Let α be a nef cohomology class on a smooth projective variety X . Then there exists an almost holomorphic, dominant rational map $f : X \dashrightarrow Y$ with connected fibers, such that*

1. α is numerically trivial on all compact fibers F of f with $\dim F = \dim X - \dim Y$
2. for every general point $x \in X$ and every irreducible curve C passing through x with $\dim f(C) > 0$, we have $\alpha \cdot C > 0$.

The map f is unique up to birational equivalence of Y .

In particular, if two general points of X can be joined by a chain C of curves such that $\alpha \cdot C = 0$, then $\alpha \equiv 0$.

2.3. The nef dimension. Since Y is unique up to a birational map, its dimension $\dim Y$ is an invariant of L which we compare to the other known invariants.

Definition 2.7. *The dimension $\dim Y$ is called the nef dimension of L . We write $n(L) := \dim Y$.*

As usual we let $\nu(L)$ be the numerical Kodaira dimension of L , i.e. the maximal number k such that $L^k \not\equiv 0$. Alternatively, if H is a fixed ample line bundle, then $\nu(L)$ is the maximal number k such that $L^k \cdot H^{n-k} \neq 0$.

Proposition 2.8. *The nef dimension is never smaller than the numerical Kodaira dimension:*

$$\nu(L) \leq n(L).$$

Proof. Fix a very ample line bundle $H \in \text{Pic}(X)$ and set $\nu := \nu(L)$. Let Z be a general member cut out by $n - \nu$ elements of $|H|$. The dimension of Z will thus be $\dim Z = \nu$, and since $L^\nu \cdot H^{n-\nu} > 0$, the restriction $L|_Z$ is big (and nef). Consequence: $\dim f(Z) = \nu$, since otherwise Z would be covered by curves C which are contained in general fibers of f , so that $L \cdot C = 0$, contradicting the bigness of $L|_Z$. In particular, we have $\dim Y \geq \dim f(Z) = \nu$ and our claim is shown. $\square$

Corollary 2.9. *The nef dimension is never smaller than the Kodaira dimension:*

$$\kappa(L) \leq n(L).$$

$\square$

2.4. The structure of the reduction map. Again let L be a nef line bundle on a projective manifold and $f : X \rightarrow Y$ a reduction map. In this section we will shed light on the structure of the map in a few simple cases.

At the present time, however, we cannot say much about f in good generality. The following natural question is, to our knowledge, open.

Question 2.10. *Are there any reasonable circumstances under which the reduction map can be taken holomorphic? Is it possible to construct an example where the reduction map cannot be chosen to be holomorphic?*

Of course, the abundance conjecture implies that the f can be chosen holomorphic in the case where $L = K_X$. Likewise, we expect a holomorphic reduction map when $L = -K_X$.

2.4.1. The case where L is big. If L is big and nef, then $n(L) = \dim X$. The converse however is false: there are examples of surfaces X carrying a line bundle L such that $L \cdot C > 0$ for all curves C but $L^2 = 0$. So in that case $n(L) = 2$, but $\nu(L) = 1$. The first explicit example is due to Mumford, see [Ha70]. This also shows that equality can fail in proposition 2.8.

2.4.2. The case where $n(L) = \dim X = 2$. In this case, we have $L \cdot C > 0$ for all but an at most countable number of irreducible curves $C \subset X$. These curves necessarily have semi-negative self-intersection $C^2 \leq 0$.

2.4.3. The case where $n(X) = 1$ and $\dim X = 2$. Here we can choose Y to be a smooth curve. The almost-holomorphic reduction $f : X \rightarrow Y$ will thus be holomorphic. In this situation, we claim that the numerical class of a suitable multiple of L comes from downstairs:

Proposition 2.11. *If $n(X) = 1$, then there exists a $\mathbb{Q}$ -divisor A on Y such that*

$$L \equiv f^*(A).$$

Proof. Since L is nef, it carries a singular metric with positive curvature current T , in particular $c_1(L) = [T]$. Since $L \cdot F = 0$, and since every $(1, 1)$ -form φ on Y is closed, we conclude that

$$f_*(T)(\varphi) = T(f^*(\varphi)) = [T] \cdot [f^*(\varphi)] = 0,$$

since $[f^*(\varphi)] = \lambda[F]$ for a fiber F of f .

Hence $f_*(T) = 0$. Let Y_0 be the maximal open subset of Y over which f is a submersion and let $X_0 = f^{-1}(Y_0)$. Then by [HL83, (18)]

$$T|_{X_0} = \sum \mu_i F_i,$$

where F_i are fibers of $f|_{X_0}$ and $\mu_i > 0$. Hence $\text{supp}(T - \sum \mu_i F_i) \subset X \setminus X_0$, an analytic set of dimension 1. Hence by a classical theorem (see e.g. [HL83]),

$$T - \sum \mu_i F_i = \sum \lambda_j G_j$$

where G_j are irreducible components of $X \setminus X_0$, i.e. of fibers, of dimension 1 and where $\lambda_j > 0$. Since $(\sum \lambda_j G_j)^2 = 0$, Zariski's lemma shows that $\sum \lambda_j G_j$ is a multiple of fibers, whence $L \equiv f^*(A)$ for a suitable $\mathbb{Q}$ -divisor A on B .

Here is an algebraic proof, which however does not easily extend to higher dimensions as the previous (see 2.3.4).

Fix an ample line bundle A on X and choose a positive rational number m such that $L \cdot A = mF \cdot A$. Let $D = mF$. Then $L^2 = L \cdot D = D^2$ and $L \cdot A = D \cdot A$. Introducing $r = L^2$ and $d = L \cdot A$, the ample line bundles $L' = L + A$ and $D' = D + A$ fulfill the equation

$$(L' \cdot D')^2 = (r + 2d + A^2)^2 = (L')^2 (D')^2,$$

hence L and D are numerically proportional, and so do L and D . Consequently $L \equiv f^*(A)$ as before. $\square$

2.4.4. The case when $n(L) = 1$ and $\dim X$ is arbitrary. The considerations of section 2.4.3 easily generalize to higher dimensions: if X is a projective manifold and L a nef line bundle on X with $n(L) = 1$, then the reduction map $f : X \rightarrow Y$ is holomorphic and there exists a $\mathbb{Q}$ -divisor on Y such that $L \equiv f^*(A)$.

2.4.5. The case where $n(L) = 0$. We have $n(L) = 0$ if and only if $L \equiv 0$.

3. A COUNTEREXAMPLE

In [Ts00], H. Tsuji claims the following:

Claim 3.1. *Let (L, h) be a line bundle with a singular hermitian metric on a smooth projective variety X such that the curvature current Θ_h is non-negative. Then there exists a (up to birational equivalence) unique rational fibration*

$$f : X \dashrightarrow Y$$

such that

- (a) *f is regular over the generic point of Y ,*

- (b) $(L, h)|_F$ is well defined and numerically trivial for every very general fiber F , and
- (c) $\dim Y$ is minimal among such fibrations.

Tsuji calls a pair (L, h) numerically trivial if for every irreducible curve C on X , which is not contained in the singular locus of h ,

$$(L, h).C = 0$$

holds. The intersection number has the following

Definition 3.2. *Let (L, h) be a line bundle with a singular hermitian metric on a smooth projective variety X such that the curvature current Θ_h is non-negative. Let C be an irreducible curve on X such that the natural morphism $\mathcal{I}(h^m) \otimes \mathcal{O}_C \rightarrow \mathcal{O}_C$ is an isomorphism at the generic point of C for every $m \geq 0$. Then*

$$(L, h).C := \limsup_{m \rightarrow \infty} \frac{\dim H^0(C, \mathcal{O}_C(mL) \otimes \mathcal{I}(h^m)/\text{tor})}{m},$$

where tor denotes the torsion part of $\mathcal{O}_C(mL) \otimes \mathcal{I}(h^m)$.

Of course, claim 3.1 is trivial as stated —except for the uniqueness assertion. What is meant is that every curve C with $(L, h).C = 0$ is contracted by f . This is also clear from Tsuji's construction.

In order to make sense of claim 3.1, statement (c) must be given the following meaning: if $x \in X$ is very general, then $(L, h).C > 0$ for all curves C through x .

There is an easy counterexample to the Claim 3.1: Take $X = \mathbb{P}^2$ with homogeneous coordinates $(z_0 : z_1 : z_2)$, $L = \mathcal{O}(1)$ and let h be induced by the incomplete linear system of lines passing through $(1 : 0 : 0)$. Then the corresponding plurisubharmonic function ϕ is given by $\frac{1}{2} \log(|z_1|^2 + |z_2|^2)$.

By [De00, 5.9], it is possible to reduce the calculation of $\mathcal{I}(h^m)$ to an algebraic problem: Since the ideal sheaf $\mathcal{J}_p$ generated by z_1, z_2 describes the reduced point $p = (1 : 0 : 0)$, it is integrally closed. Let $\mu : \mathbb{F}_1 \rightarrow \mathbb{P}^2$ be the blow up of $\mathbb{P}^2$ in p . Then $\mu^*\mathcal{J}_p$ is the invertible sheaf $\mathcal{O}(-E)$ associated with the exceptional divisor E . Now, one has $K_{\mathbb{F}_1} = \mu^*K_{\mathbb{P}^2} + E$. By the direct image formula in [De00, 5.8] it follows

$$\mathcal{I}(m\phi) = \mu_*(\mathcal{O}(K_{\mathbb{F}_1} - \mu^*K_{\mathbb{P}^2}) \otimes \mathcal{I}(m\phi \circ \mu)).$$

Now, $(z_i \circ \mu)$ are generators of the ideal $\mathcal{O}(-E)$, hence

$$m\phi \circ \mu \sim m \log g,$$

where g is a local generator of $\mathcal{O}(-E)$. But $\mathcal{I}(m\phi \circ \mu) = \mathcal{O}(-mE)$, hence

$$\mathcal{I}(m\phi) = \mu_*\mathcal{O}_{\mathbb{F}_1}((1-m)E) = \mathcal{J}_p^{m-1}.$$

Let $C \subset \mathbb{P}^2$ be a smooth curve of degree d and genus g . If $p \notin C$ then $\mathcal{O}_C(mL) \otimes \mathcal{J}_p^{m-1} = \mathcal{O}_C(mL)$, and $(L, h).C = L.C = d$. On the other hand, if $p \in C$, then

$$\mathcal{O}_C(mL) \otimes \mathcal{J}_p^{m-1} = \mathcal{O}_C(mL - (m-1)p).$$

The degree of this invertible sheaf is $md - m + 1 = m(d - 1) + 1$, hence by Riemann-Roch

$$\begin{aligned} \limsup_{m \rightarrow \infty} \frac{\dim H^0(C, \mathcal{O}_C(mL) \otimes \mathcal{I}(m\phi))}{m} &= \limsup_{m \rightarrow \infty} \frac{m(d - 1) + 1 + 1 - g}{m} \\ &= d - 1. \end{aligned}$$

Consequently, Tsuji's fibration could only be the trivial map $\mathbb{P}^2 \rightarrow \mathbb{P}^2$. It can't be the map to a point, because there are curves with intersection number ≥ 1 on $\mathbb{P}^2$, and it can't be an almost holomorphic rational map to a curve, because $\mathbb{P}^2$ does not contain curves with vanishing self-intersection. This contradicts condition (c) in the claim.

It remains an open question if claim 3.1 is true without condition (a), and if in the case of singular hermitian metrics induced by linear systems, the fibration may be taken as the induced rational map.

REFERENCES

- [Ca81] F. Campana. Coréduction algébrique d'un espace analytique faiblement kählérien compact. *Inv. Math.*, 63:187-223, 1981.
- [Ca94] F. Campana. Remarques sur le revêtement universel des variétés kähleriennes compactes. *Bull. SMF*, 122:255-284, 1994
- [De00] J.P. Demailly. Vanishing theorems and effective results in algebraic geometry. Trieste lecture notes, 2000
- [Ha70] R. Hartshorne. *Ample subvarieties of algebraic varieties*, volume 156 of *Lecture Notes in Mathematics*, Springer, 1970.
- [HL83] R. Harvey, H.B. Lawson. An intrinsic characterization of Kähler manifolds. *Inv. math.*, 74:169-198, 1983.
- [Pa98] M. Paun. Sur l'effectivité numérique des images inverses de fibrés en droites. *Math. Ann.*, 310:411-421, 1998.
- [Ts00] H. Tsuji. Numerically trivial fibrations. LANL-preprint math.AG/0001023, October 2000.

THOMAS BAUER, THOMAS ECKL, STEFAN KEBEKUS AND THOMAS PETERNELL: INSTITUT FÜR MATHEMATIK, UNIVERSITÄT BAYREUTH, 95440 BAYREUTH, GERMANY

E-mail address: thomas.bauer@uni-bayreuth.de

E-mail address: thomas.eckl@uni-bayreuth.de

E-mail address: stefan.kebekus@uni-bayreuth.de

E-mail address: thomas.peternell@uni-bayreuth.de

FRÉDÉRIC CAMPANA: DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ NANCY 1, BP 239, 54507 VANDOEUVRE-LES-NANCE CÉDEX, FRANCE

E-mail address: campana@iecn.u-nancy.fr

SLAWOMIR RAMS: MATHEMATISCHES INSTITUT DER UNIVERSITÄT, BISMARCK-STRASSE 1½, 91054 ERLANGEN, GERMANY

E-mail address: rams@mi.uni-erlangen.de

TOMASZ SZEMBERG: UNIVERSITÄT GH ESSEN, FACHBEREICH 6 MATHEMATIK, 45117 ESSEN, GERMANY

E-mail address: mat905@uni-essen.de

LORENZ WOTZLAW: MATHEMATISCHES INSTITUT, HUMBOLDT-UNIVERSITÄT BERLIN,
10099 BERLIN, GERMANY

E-mail address: `wotzlaw@mathematik.hu-berlin.de`